



*Original Article*

## COMPARISON BETWEEN ETHANOLIC EXTRACTS OF THE PLANT MORINGA PTERYGOSPERMA, AGAINST CARBON TETRACHLORIDE INDUCED HEPATOPATHY

Prabhat Kumar Das,<sup>1\*</sup> Dr. Prasanna Panda,<sup>2</sup> Somya Ranjan Pani,<sup>2</sup> Ranjan Sethi<sup>1</sup>

**Affiliated to:**

<sup>1</sup> GRY Institute of Pharmacy, Borawan, Khargome, MP-451228

<sup>2</sup> University Department of Pharmaceutical Sciences, Utkal University, Orissa.

### ABSTRACT

**Object:** Comparison between protective effects of the ethanolic extracts of the plant Moringa pterigosperma against CCl<sub>4</sub> induced hepatic failure in male albino rats was investigated. **Materials and methods:** For acute and massive invasion of hepatopathy, CCl<sub>4</sub> (i.p injection of CCl<sub>4</sub>+Olive Oil in 1:1 ratio; 2ml/kg) was used and the insidious intoxication was evidenced by evaluation of various biochemical parameters followed by significant weight loss in toxic control group. The administration of ethanolic extracts (250mg/kg and 150mg/kg of body weight) for 7 days, elicited protective action since the elevated levels of marker enzymes (AST, ALT, ALP) of liver functions were found to be decreasing progressively in a dose dependent manner with net weight gain. **Results:** In the ethanolic extract 250mg/kg treated rat group all the marker enzymes were analyzed to be decreasing significantly and the final body weight was also significantly increased when compared with the toxic control group. The serum total protein and the serum albumin were also approaching normal values. The results found in ethanolic extract 250mg/kg treated rat were quite promising and were comparable with a standard polyherbal drug Liv-52. **Conclusion:** The statistically processed results support the conclusion, that the ethanolic extracts of Moringa pterigosperma whole plant (250mg/kg and 150mg/kg) possesses dose dependent, significant protective activity against CCl<sub>4</sub> induced hepatotoxicity.

**Keywords:** Hepatotoxicity, marker enzymes, Asparate transaminase, Alanine aminotransferase, Alkaline Phosphatase.

\*Corresponding Author:

Prabhat Kumar Das

GRY Institute of Pharmacy, Borawan, Khargome,  
MP-451228

E Mail: suchiprabhat3027@yahoo.co.in



## 1.0 INTRODUCTION <sup>[1] [2][3] [4][5][6]</sup>

The aim of the present context is to evaluate the hepatoprotective potential of the ethanolic extract of *Moringa pterygosperma* whole plant in a conventional animal model of hepatopatotoxicity.

*Moringa pterygosperma* belongs to a onogeneric family of shrubs and tree, Moringaceae and is considered to have its origin in Agra and Oudh, in the northwest region of India, south of the Himalayan Mountains. They are an exceptionally good source of provitamin A, vitamins B, and C, minerals (in particular iron), and the sulphur-containing amino acids methionine and cystine. The seed oil of *Moringa* contains approximately 13 % saturated fatty acids and 82 % unsaturated fatty acids. It has a particularly high level of oleic acid (70 %). *Moringa* leaves contain more Vitamin A than carrots, more calcium than milk, more iron than spinach, more Vitamin C than oranges, and more potassium than bananas, and that the protein quality of *Moringa* leaves rivals that of milk and eggs. The seeds contain 19 to 47 percent oil Known commercially as “ben oil”, it is similar to olive oil and is rich in palmetic, stearic, behmic, and oleic acids, and are used for human consumption, and in cosmetics and soaps. In traditional Indian medicine various parts of the tree are used therapeutically, including for treatment of ascites, rheumatism, venomous bites,

and as cardiac, anti diabetic and circulatory stimulants. The root of young trees and also the root bark are considered rubefacient, vescicant carminative, stomachic, and abortifacient; among other uses, they are commonly applied externally to cure inflammatory swellings. The flowers and roots contain pterogosperrin, an antibiotic that is highly effective in the treatment of cholera. The flowers are also used as a tonic and diuretic. The leaves, rich in vitamin A and C, are considered useful in scurvy and respiratory ailments; they are also used as an emetic. The juice extracted from the leaves has strong antibacterial and antimalarial properties a paste of the leaves is used as an external application to promote healing of wounds. The seed oil is applied externally to treat rheumatism and gout.

$\text{CCl}_4$  is a widely used industrial chemical and a potent hepatotoxin. It induces hepatotoxicity by producing free radical, putting oxidative stress hence causing lipid peroxidation in liver tissues, consequently necrotic liver damage <sup>[7]</sup>.

## 2.0 MATERIALS AND METHODS

### 2.1 Plant material

The whole part of plant *Moringa pterygosperma* plant was collected from young matured plant from the rural belt of Balasore, Orissa during the month of Jan-Feb and identified by the botanist of Department of Botany, Utkal University, Bhubaneswar by comparing with the voucher specimen present in the herbarium. After



authentication fresh plant materials were collected in bulk, washed under running tap water to remove adhering dust, dried under shade and pulverized in a mechanical grinder. The coarse powder was used for further studies.

#### Animals

Healthy albino male rats of Wistar strain weighing between 150-200 g were selected for the investigation (Brought from Ghosh Enterprises, Kolkata, India). The animals were kept under maintained uniform laboratory conditions ( $12 \pm 1$  hr, day and night Schedule; temperature maintained between  $11-20^\circ \pm 2^\circ\text{C}$ ; housed in large spacious hygienic cages) before a week of the experiment for acclimatization. The approval of the protocol by I.A.E.C/U.D.P.S/990/2005 -Vanivihar, Bhubaneswar. Following C.P.C.S.E.A guideline.

#### 2.2 Chemicals

Chemicals used in the study were of analytical grade and were procured from Merck specialties private limited, Mumbai, India. All biochemical assay kits were purchased from Ecoline: Merck specialties private limited, Mumbai, India.

#### 2.3 Extraction of plant material and preparation of test dose

About 200 gm of coarse dried powder of whole plant of the *Moringa pterygosperma* was taken in the soxhlet apparatus and extracted successively<sup>[8]</sup> using the selected solvents in order (i.e. Pet. ether→Chloroform→Ethanol). The extraction for each solvent was carried out for 18

to 20 hours. The extract was collected by evaporating the solvents by slow heat treatment. Total 1.4kg of pulverized stem was subjected under solvent extraction to produce the required amount of test extract.

Calculated amount of dried ethanolic extract was suspended in 0.5% w/v of sodium-CMC in normal saline solution to get the test doses (200mg/kg and 400mg/kg per ml.). The dose limits were selected on the basis of previously performed oral acute toxicity studies in mice, in accordance with the OECD guidelines.<sup>[9]</sup>

#### 2.4 Acute toxicity studies:

In the acute toxicity test carried out in mice we take six doses and 10 mice in each dose of both aqueous and ethanolic extract i.e. 500, 1000, 1500, 2000, 2500, 3000 mg/kg body weight. All groups of test drug showed neither any toxic effect nor any lethal effect in the dose range of 500 to 3000 mg/kg body weight. In 3000 mg/kg of ethanolic extract showed altered behavior of some mice had been observed. So we had taken a minimum and maximum dose 150mg/kg and 250 mg/kg of body weight for ethanolic extract for further screenings.

#### 2.5 Hepatoprotective activity study:

30 rats were divided randomly into 5 groups, each comprising 6 animals. Group I (Normal Control) received oral dose of 0.5% Sodium CMC (1 ml each) for 7 days. Group II (Toxic Control) received single dose of  $\text{CCl}_4$  ( $\text{CCl}_4$  + Olive oil in 1:1 ratio; 2ml/kg of body wt; i.p.) on day 1 and day 7 of the experiment<sup>[10]</sup>. Group III, IV and V received standard polyherbal





drug 'Liv-52' (5 ml/kg; p.o.) <sup>[11]</sup>(Liv-52 Syrup, Himalaya Drug Company., Bangalore, India), ethanolic extract 250 mg/kg of body wt. and 150mg/kg of body wt. once in a day for 7 days respectively, along with the i.p dose of CCl<sub>4</sub> on day 1 and day 7 as mentioned above. The treatment duration was of 7 days. On the 8<sup>th</sup> day of the study the animals were sacrificed under anesthesia and blood samples were collected from each animal to produce the serum for biochemical assay. The biochemical investigations were performed by using a Biochemical semi auto analyzer (EBRA-Chem-5 Plus. V2., West-Germany). The biochemical parameters considered were: Serum AST (SGOT) i.e. Asparate transaminase, ALT (SGPT) i.e. Alanine aminotransferase, ALP (alkaline phosphatase), Serum albumin and total protein. <sup>[12][13][14]</sup>

## 2.6 Statistical Analysis.

Table 1: Change in body weight, Serum albumin, total protein and assay of Marker Enzymes on 8<sup>th</sup> day.

Group.	Change in Body wt. (gm)	Albumin. (gm/dl)	Total protein. (gm/dl)	AST. (IU/l)	ALT. (IU/l)	ALP. (IU/l)
Normal Control.	10.66±1.28	2.83±0.17	5.70±0.35	75.04±7.23	49.21±4.90	80.30±3.58
Toxic Control.	-12.83±1.13 <sup>c</sup>	1.40±0.10 <sup>c</sup>	3.11±0.16 <sup>c</sup>	840.37±28.26 <sup>c</sup>	746.75±18.94 <sup>c</sup>	133.65±3.47 <sup>c</sup>
Standard.	6.83±0.70 <sup>c</sup>	2.71±0.27 <sup>c</sup>	5.09±0.20 <sup>c</sup>	246.23±16.93 <sup>c</sup>	186.38±14.60 <sup>c</sup>	86.49±7.17 <sup>c</sup>
EEMP <sub>150</sub> Group.	3.33±1.90 <sup>c</sup>	2.55±0.16 <sup>b</sup>	4.70±0.36 <sup>b</sup>	393.08±12.29 <sup>c</sup>	241.07±11.73 <sup>c</sup>	112.58±6.82
EEMP <sub>250</sub> Group.	6.16±1.01 <sup>c</sup>	2.61±0.18 <sup>c</sup>	4.96±0.31 <sup>b</sup>	272.77±24.08 <sup>c</sup>	189.15±7.16 <sup>c</sup>	97.15±6.54 <sup>b</sup>

a - p<0.05, b - p<0.01, c - p<0.001. For n=6. Toxic control group was compared with Normal control group and all the other groups were compared with toxic control group.

- EEMP= Ethanolic extract of the plant Moringa pterygosperma
- AST = Asparate transaminase.

- ALT = Alanine aminotransferase.
- ALP = Alkaline Phosphatase.

Histogram no.1:


Results of all the estimations done were indicated in terms of Mean ± SEM. Statistical significance of data were assessed by analysis of variance (One Way- ANOVA), followed by comparison between different groups using 'Tukey-Kramer' multiple comparison test. The significance was set at the level of p<0.05. The toxic control group was compared with the normal control group and all other treatment groups were compared with the toxic control group.

## 3.0 RESULT AND DISCUSSION

The biochemical assays of hepatoprotective study were performed on 8<sup>th</sup> day. The results were presented in Table 1. (Change in body weight; Serum albumin and total protein) and Table 2. (Assay of marker enzymes).




**Graphical representation of the Assay of Marker Enzymes on 8th day**



Histogram no.2:

**comparison of serum albumine and total protein of different group in CCl<sub>4</sub> induced hepatopathy on 8th day**



- 1 = Normal control group
- 2 = Toxic control group
- 3 = standard group
- 4 = Ethanolic extract of the plant *Moringa pterygosperma* 150 mg/kg treated rats
- 5 = Ethanolic extract of the plant *Moringa pterygosperma* 250 mg/kg treated rats

A significant elevation in serum AST, serum ALT, serum ALP levels and significant decrease in serum Total Protein and serum albumin level followed by significant weight loss were found in Toxic control group, when compared with the Normal control group.

In the EEMP<sub>150</sub> group (ethanolic extract of the plant 150mg/kg treated rat group), the final body wt. was observed to be elevated as the change in body wt. which was found to be significant. Both the serum AST and the serum ALT were found to be significantly low when compared with the



Toxic control group. But the serum ALP level was not significantly decreased. The serum albumin and serum total protein were likely to be approaching normal values and found significant 1, when compared with that of Toxic control group.

The EEMP<sub>250</sub> group (ethanolic extract of the plant 250mg/kg treated rat group), the final body wt. was significantly elevated, when compared with the Toxic control group. The serum total protein and the serum albumin were almost approaching normal values and comparable with the results observed in Standard group. Serum AST and serum ALT were found to be significantly lower and serum ALP level was also significantly decreased (when compared with the Toxic control group), approaching almost normal values as that of observed in case of Standard group, supports the restoration of normal hepatic functions.

The above results suggest that ethanolic extract treated rats (at a dose level of 250mg/kg and 150mg/kg treated rats) had gained normalcy against the hepatocellular injury caused by CCl<sub>4</sub> during the 7 day treatment period and both dose levels were found almost equipotent. The result of these investigations was comparable and matches the previously reported protective effects of other plants.<sup>[15][16]</sup>

#### 4.0 CONCLUSION

In the present pharmacological evaluation the whole plant extracts (ethanolic) of *Moringa pterygosperma* plant was extensively investigated for its Hepatoprotective potential against substance (CCl<sub>4</sub>) induced hepatopathy. At the end of our study, a strong conclusion can be drawn that, the ethanolic extract of *Moringa pterygosperma* possess Hepatoprotective activities more or less depending on the dose levels. The ethanolic extract of the plant at a dose level of 250mg/kg exhibited competent, potent and comparable results where as ethanolic extract at a dose level of 150mg/kg, observed to have moderate level of efficacy, promoting *Moringa pterygosperma* plant as a promising Hepatoprotective plant species, seeking vast multidimensional future research work up to the molecular level to establish new up-to-date scientific data about this plant species and to elucidate its exact mechanism of protective effect. Future studies may be aimed at hepatoprotective study in other chronic models of Hepatopathies, antioxidant and free radical scavenging potentials, toxicological studies and other pharmacological activities as well.

#### ACKNOWLEDGEMENTS

The authors are very much grateful to the Sigma Aldrich Pvt. Ltd., Hyderabad branch and Bangalore branch for their kind support to carry out this work in time. Thanks to Professor B.B.Barik, HOD of University Department of





Pharmaceutical Sciences, Utkal University and Professor P.K.Panda, HOD of Pharmacology dept., for providing necessary lab facilities and kind co- operation to carry out this work in time. I also give my heartfelt thanks to Dr. Debasish Pradhan and Dr. S. Sahoo to suggest this topic and give their important suggestions and help me time to time during my project work.

## REFERENCE

1. Foidl N.et al., The potential of Moringa oleifera for agricultural and industrial uses, October 20th - November 2nd 2001.
2. Jed W. Fahey, Moringa oleifera: A Review of the Medical Evidence for Its Nutritional, Therapeutic, and Prophylactic Properties. Part 1, review, Trees for Life Journal a forum on beneficial trees and plants
3. Kar A, B Choundhary, and N Bandyopadhyay (2003) Comparative evaluation of hypoglycaemic activity of some Indian medicinal plants in alloxan diabetic rats. Journal of Ethnopharmacology 84(1): 105-108.
4. Limaye DA, AY Nimbkar, R Jain, and M Ahmad (1995) Cardiovascular effects of the aqueous extract of Moringa pterygosperma. Phytotherapy Research 9: 37-40.
5. Pal SK, PK Mukherjee, and BP Saha (1995) Studies on the antiulcer activity of Moringa oleifera leaf extract on gastric ulcer models in rats. Phytotherapy Research 9: 463-465.
6. Pari L and NA Kumar (2002) Hepatoprotective activity of Moringa oleifera on antitubercular drug-induced liver damage in rats. Journal of Medical Foods 5(3): 171-177.
7. Ram VJ, Goel A (1999) Curr. Med. Chem., 6, 217-254
8. Suftness M, Douras J (1979) Drugs of Plant origin in methods in cancer research, Academic press, New York, 548 pp
9. OECD (2000) Acute oral Toxicity - Acute oral toxic class method, Guideline 423, adopted 23.03.1996. In: Eleventh Addendum to the OECD guidelines for the Testing of Chemicals. Organization for Economic Co-Operation and Development, Paris.
10. Ray D, Sharatchandra K, Thokchom IS (2006) Antipyretic, antidiarrhoeal, hypoglycemic and hepatoprotective activities of ethyl acetate extract of Acacia catechu wild. in albino rats. Ind. Jonl. Pharmacology, 38 (6), 408-413
11. Sandhir R, Gill KD (1999) Hepatoprotective Effects of Liv.52 on Ethanol-induced Liver Damage in Rats. Indian Journal of Experimental Biology, 37, 762-766
12. Cheesbrough M (2003) District Laboratory Practice in Tropical Countries, Cambridge University Press, 310-395 pp
13. Tietz (1999) Text Book of Clinical Chemistry (3<sup>rd</sup> Edn), W.B Saunders Company, Philadelphia, 617-721 pp
14. Doumas BT, Watson WA (1971) J. Clin. Chem., 31, 87



15. Dwivedi Y, Rastogi R, Chander R, Sharma SK, Kapoor NK, Garg NK (1990) Hepatoprotective activity of Picroliv against carbontetrachloride induced liver damage in rats. Indian J Med Res, 92, 185-200
16. Mohamed AF, Hasan AGA, Hamany MI, Sattar EA (2005) Antioxidant and Hepatoprotective effects of Eucalypts maculate. Med. Sci. Monit., 11 (11), 426-431.