



ESTROGENIC ACTIVITY OF HYDRO-ALCOHOLIC EXTRACT OF *ARTOCARPUS LAKOOCHA* FRUIT ON FEMALE SPRAGUE-DAWLEY RATS

Jawaid Talha¹, Gautam Piyush¹, Rahman Md. Azizur², Kamal Mehnaz^{2,*}

¹Department of Pharmacology, Hygia Institute of Pharmaceutical Education and Research, Lucknow, UP (India) 226020

²Faculty of Pharmacy, Integral University, Lucknow, U.P. (India) 226026

ARTICLE INFO	ABSTRACT
<p>Published on: 15-6-2015 ISSN: 0975-8216</p> <hr/> <p>Keywords: <i>Artocarpus lakoocha</i>; estrogenic activity; uterotropic assay; vaginal cytology; vaginal opening</p> <hr/> <p>Corresponding author: Dr. Mehnaz Kamal, Faculty of Pharmacy, Integral University, Lucknow, U.P. (India) 226026</p>	<p>Aim: Hormone replacement therapy was considered a safe treatment for short-term therapy. But, because of adverse outcomes such as increased risk of endometrial and breast cancer, stroke and pulmonary thrombo-embolism after its long-term use, women turned to use natural health remedies. Thus, the estrogenic activity of 70% hydro-alcoholic extract of <i>Artocarpus lakoocha</i> fruit (HEAL) was evaluated.</p> <p>Materials and Methods: HEAL was subjected to phytochemical analysis as per standard protocol. Estrogenic potential of HEAL was performed by uterotropic assay, vaginal cytology and measurement of vaginal opening at different doses of 200 mg/kg, 400 mg/kg and 600 mg/kg bw <i>po</i> in Sprague-Dawley (SD) rats and compared with standard drug β-estradiol (2 mg/kg bw, <i>po</i>). Ovariectomized immature and mature female SD rats were administered with HEAL for 7 days.</p> <p>Results: It showed the presence of tannins, saponins, triterpenes, flavonoids, steroids and alkaloids as major secondary metabolites in HEAL. Dose of 600 mg/kg administered in ovariectomized immature and mature female SD rats resulted in significant increase in the uterine wet weight of 219.60\pm8.71 mg and 906.19\pm26.91 mg respectively when compared with that of control rats of 110.55\pm1.16 mg and 505.66\pm27.38 mg respectively. Only cornified epithelial cells were seen in HEAL-treated rats at the dose of 600 mg/kg indicating the presence of estrogen. Significant vaginal opening was also seen in HEAL-treated rats at the dose of 600 mg/kg.</p> <p>Conclusion: It was concluded that HEAL possess promising estrogenic activity.</p>

INTRODUCTION

Estrogen plays an extremely essential role in growth regulation and function in numerous female target organs such as vagina, uterus,

skeletal and cardiovascular systems^[1,2].

Estrogen deficiency results in cessation of ovarian production which is associated with many complaints experienced by menopausal

women^[3,4,5]. For decades, the hormone replacement therapy (HRT) had been used as an alternative for managing menopause-induced complaints^[6,7]. It is presumably a safe treatment for short-term therapy. But, it was implicated in adverse outcomes such as increased risk of endometrial and breast cancer, stroke and pulmonary thrombo-embolism after long-term use^[8,9]. Due to this, women turned to use natural health remedies. Thus, the interest in plant-derived phyto-estrogens raised significantly in the last decades.

The genus *Artocarpus* comprises approximately 50 species of deciduous and evergreen trees. It is economically important as a source of edible fruit and yields good timber. It is widely used in traditional folk medicines in South-East Asia and India for the treatment of several diseases like abscess, ulcer, inflammation, diarrhoea, diabetes, malaria and tapeworm infection^[10,11,12]. *Artocarpus lakoocha* fruits are eaten fresh. Its edible fruit pulp act as a liver tonic. Its male flower spikes and raw fruits are used in pickles and chutney. Its brown powder named as Puag-Haad in Thailand is used as a traditional anthelmintic drug for the treatment of tapeworm infection^[13,14]. Its bark containing tannin is chewed like betel nuts and is also used to treat skin diseases^[15]. This plant has antiviral and anthelmintic activities and has also been used as skin-whitening agent in cosmetics^[16,17,18]. *Artocarpus lakoocha* plant contains various phytoconstituents like β -sitosterol, cycloartenol, cycloartenone, α -amyrin acetate and lupeol acetate. It also contains artocarpin, norartocarpin, cycloartocarpin, resorcinol, oxyresveratrol, pentosans, lignin, tannin, holocellulose and α -

cellulose^[19]. The aim of study was designed for exploration purpose to evaluate the estrogenic activity of HEAL in SD rats by uterotrophic assay, vaginal cytology and measurement of vaginal opening.

MATERIALS AND METHODS

COLLECTION AND AUTHENTICATION OF PLANT SPECIMEN

The fruits of *Artocarpus lakoocha* used in the present study were collected from local market of Lucknow, India and authenticated by National Botanical Research Institute (NBRI), Lucknow (authentication reference number NBRI/CIF/260/2011).

PREPARATION OF HYDRO-ALCOHOLIC EXTRACT

Fruits were cut into small pieces and shade dried. The dried material was then powdered into coarse powder by a mechanical grinder. The dried coarse powder was evenly packed in a Soxhlet extractor for extraction. It was defatted with petroleum ether (60-80°). The temperature was maintained on an electric heating mantle with thermostat control. The defatted marc was further extracted with 70% ethanol for 24 hr. Extract was filtered and then concentrated to dryness in a rotavapor under reduced pressure and controlled temperature. It was stored in refrigerator and kept in desiccator few hours before use^[20].

MATERIALS AND REAGENTS

β -Estradiol (National chemicals, Vadodara, Gujrat), methanol (Ranbaxy, India), normal saline (NaCl 0.9%), methylene blue, ketamine hydrochloride injection (Neon Laboratories Limited), chloroform (RFCL Ltd, New Delhi).

PRELIMINARY PHYTOCHEMICAL SCREENING

Phytochemical screening of HEAL was carried out for alkaloids (Mayer test), saponins (Foam test), flavonoids (Shinoda test), carbohydrates (Benedict test), tannins (Ferric chloride test) steroids and triterpenes (Lieberman-Burchard test) according to standard methods.^[21]

ANIMALS

Immature and mature female SD rats weighing between (60-70±10, 150-200±20 gm respectively) procured from the Central Drug Research Institute Lucknow, Uttar Pradesh were housed in polypropylene cages (22.5×37.5 cm²) and maintained under standard laboratory environmental conditions viz., temperature 25±2°C, 12 h light:12 h dark cycle and 55±10% relative humidity with free access to standard pellet diet and water, *ad libitum*. The experimental protocol was approved by the Institutional Animal Ethics Committee, which follow the guidelines of the Committee for the Purpose of Control and Supervision of Experiments on Animals (CPCSEA) and conform to the international norms of the Indian National Science Academy. Ethical norms were strictly followed during all experimental procedures [Hygia/M.Pharm./05/2011-12].

PREPARATION OF TEST SAMPLES AND DOSING

The ovariectomized rats were divided into 5 groups each consisting of 5 animals. Group 1 (control) received 0.6% (w/v) sodium CMC suspension (10 ml/kg, *po*), Group 2 (standard) aqueous suspension of β -estradiol in 0.6% (w/v) sodium CMC (2 mg/kg, *po*), Group 3 (test) aqueous suspension of HEAL in 0.6%

(w/v) sodium CMC (200 mg/kg, *po*), Group 4 (test) aqueous suspension of HEAL in 0.6% (w/v) sodium CMC (400 mg/kg, *po*) and Group 5 (test) aqueous suspension of HEAL in 0.6% (w/v) sodium CMC (600 mg/kg, *po*).

MEASUREMENT OF ESTROGENIC PARAMETERS

UTEROTROPIC ASSAY IN IMMATURE AND MATURE OVARIECTOMIZED FEMALE RATS

Immature and mature female SD rats were divided into five groups, each of five rats. All rats were bilaterally ovariectomized by dorso-lateral approach under anesthesia and semi-sterile conditions. After one day of ovariectomization, the dose treatment was started. All the above mentioned dosages were administered orally daily for 7 days. During this period of treatment, the rats were maintained under standard pellet diet and water. Vaginal cornification was examined daily. After 24 h of last treatment, hysterectomy was performed in all rats under ketamine anesthesia. Harvested uteri were cleaned carefully from adhering connective tissue and weighed.^[22]

VAGINAL CYTOLOGY

This was performed by vaginal smear method in ovariectomized immature female SD rats. Vaginal smear was taken by introducing a few drops of saline into vagina with the help of eye dropper. The saline was expelled into the vagina and withdrawn two or three times. The contents of the eye dropper was placed and spread on a glass slide, the smear was immediately fixed with 1% w/v aqueous methylene blue for 5-6 min. The smear was examined under microscope to check the

presence or absence of leukocytes and cornified epithelial cells. The presence of only cornified epithelial cells is indicative of estrogenic activity.^[23]

The vaginal smear test score report was tested.
 0 Di estrus smear, mainly leucocytes few epithelial cells.

1 Mixture of leucocytes and epithelial cells.

2 Pro estrus smear nucleated or nucleated+cornified cells.

3 Estrus smear, cornified cells only.

MEASUREMENT OF VAGINAL OPENING

The vaginal opening was observed and noted daily for 7 days after oral administration of HEAL in ovariectomized immature female SD rats. Increase in vaginal opening is indicative of estrogenic activity.^[24]

STATISTICAL ANALYSIS

The results were analyzed using one way analysis of variance (ANOVA) with Dunnett's

comparison test. *P* values<0.01 were considered statistically significant.

RESULTS

PRELIMINARY PHYTOCHEMICAL SCREENING

HEAL showed the presence of carbohydrates, saponins, triterpenes, tannins, flavonoids, alkaloids and steroids.

UTEROTROPIC ASSAY IN IMMATURE AND MATURE OVARIECTOMIZED FEMALE RATS

Oral administration of β -estradiol (positive control) at 2 mg/kg, and HEAL at 200, 400 and 600 mg/kg, *po*, respectively, effectively increased the uterine wet weight both in immature and mature ovariectomized rats. Increase in uterine wet weight was dose-related (**Figure 1** and **Figure 2**). HEAL at the dose of 600 mg/kg, *po* showed estrogenic activity comparable to that of standard β -estradiol.




Figure 1 Effects of HEAL 200 mg/kg, 400 mg/kg and 600 mg/kg, *po* on uterine wet weight in immature SD rat. β -estradiol was used as standard at 2 mg/kg b.w. Data were expressed as mean \pm S.E.M. ****P*<0.01 (significantly different from the control)




Figure 2 Effects of HEAL 200 mg/kg, 400 mg/kg and 600 mg/kg, *po* on uterine wet weight in mature SD rat. β -estradiol was used as standard at 2 mg/kg b.w. Data were expressed as mean \pm S.E.M. *** P <0.01 (significantly different from the control)


VAGINAL CYTOLOGY

Animals showing the score 2 and 3 were considered to be positive. The vaginal smear test score of cornification of epithelial cells was positive because the number of cornified cells in the vaginal smear was considerably higher than the control and less than β -estradiol treated animals as shown in **Table 1** and **Figure 3**. HEAL treated rats (at the dose of 600 mg/kg, *po*) showed only cornified epithelial cells. This result clearly indicated that HEAL at a dose of 600 mg/kg, *po* is having significant estrogenic activity.

Table 1 Effects of HEAL on vaginal cornification in ovariectomized immature rats

Group	Drug treatment	Dose	Vaginal smear test score
1.	Control	10 ml/kg	Negative
2.	Standard	2 mg/kg	Positive
3.	HEAL	200 mg/kg	Negative
4.	HEAL	400 mg/kg	Negative
5.	HEAL	600 mg/kg	Positive

Animals showing the score 2 and 3 are considered to be positive.






Figure 3 Photomicrograph of methylene blue stained vaginal smear (A) in diestrous of control rat, showing only leukocytes. (B) in estrous of β -estradiol (2 mg/kg, *po*) treated rat, showing only cornified epithelial cells. (C) in estrous of HEAL (200 mg/kg, *po*) treated rat, showing only leukocytes. (D) in estrous of HEAL (400 mg/kg, *po*) treated rat, showing only few cornified epithelial cells. (E) in estrous of HEAL (600 mg/kg, *po*) treated rat, showing only cornified epithelial cells

VAGINAL OPENING

HEAL showed a dose dependent increase in vaginal opening from day fourth onwards compared to control which remained closed. HEAL at dose of 600 mg/kg, *po* showed significant vaginal opening. But, HEAL at

dose of 200 mg/kg, *po* and 400 mg/kg, *po* showed insignificant vaginal opening (Table 2 and Figure 4). It showed that HEAL at dose of 600 mg/kg, *po* is having significant estrogenic activity.

Table 2 Effects of HEAL on vaginal opening in immature ovariectomized rats

Group	Drug treatment	Dose	Vaginal opening (%)						
			Day 1	Day 2	Day 3	Day 4	Day 5	Day 6	Day 7
1.	Control	10 ml/kg	0	0	0	0	0	0	0
2.	Standard	2 mg/kg	0	0	0	70	100	90	80
3.	HEAL	200 mg/kg	0	0	0	5	5	6	8
4.	HEAL	400 mg/kg	0	0	0	45	55	50	60
5.	HEAL	600 mg/kg	0	0	0	60	85	75	70




Figure 4 Effects of HEAL on vaginal opening in ovariectomized immature SD rats. (A) Control (B) Standard (C) HEAL 200mg/kg, *po* (D) HEAL 400mg/kg, *po* (E) HEAL 600mg/kg, *po*

DISCUSSION

The excess of estrogen can cause breast, endometrial, ovarian, and prostate cancer and its deficiency can result in menopausal symptoms, cardiovascular disease and

osteoporosis. The major causes of estrogen deficiency in females are menopause and ovariectomy. Interest in plant derived estrogens or phytoestrogens has recently been amplified by the realization that HRT is not as

safe or effective as previously considered.^[25,26] Phytoestrogens have negative impact on the development of human reproductive organs and male fertility, unlike xenobiotic estrogens (environmental pollutants with estrogenic activity), they are believed to have primarily beneficial effects on the health.^[27] Results of uterotrophic assay revealed that HEAL showed increase in uterine wet weight after administration of 200 mg/kg, 400 mg/kg and 600 mg/kg, *po* dose both in immature and mature ovariectomized rats. It clearly indicated that all the doses contain estrogen like compound. HEAL (600 mg/kg, *po*) treated rats showed only cornified epithelial cells. The vaginal opening showed that HEAL (600 mg/kg, *po*) is having significant estrogenic activity.

CONCLUSION

HEAL possess significant estrogenic activity at dose of 600 mg/kg, *po* which was evident by uterotrophic assay, measurement of vaginal opening and vaginal cytology in female rats. HEAL could be useful as a safe natural source for estrogenic activity for postmenopausal women.

REFERENCES

1. Couse JF, Korach SK. Estrogen receptor null mice: what have we learned and where will they lead us? *Endocrine Rev.* 1999; 20: 358-417.
2. Korach KS, Migliaccio S, Davis VL. Estrogens. In: Munson PL, Mueller RA, Breese GR (Eds.). *Principles of Pharmacology: Basic Concepts and Clinical Applications*. New York: Chapman and Hall; 1995. p. 809-825.
3. Mikkola TS, Clarkson TB. Estrogen replacement therapy, atherosclerosis, and vascularfunction. *Cardiovas Res.* 2002; 53: 605-619.
4. Turner RT, Riggs BL, Spelsberg TC. Skeletal effects of estrogen. *Endocrine Rev.* 1994; 15: 275-300.
5. Versi E, Harvey MA, Cardozo L, Brincat M, Studd JWW. Urogenital prolapsed and atrophy at menopause. A prevalence study. *Int Urogynecol J.* 2001; 12:107-110.
6. Greendale GA, Reboussin BA, Hogan P, Barnabei VM, Shumaker S, Barrett-Connor E. Symptom relief and side effects of postmenopausal hormone results from the postmenopausal estrogen/progestin interventions trial. *Obstet Gynecol.* 1998; 92:982-988.
7. Nichols KC, Schenkel L, Benson H. 17-beta-Estradiol for postmenopausal estrogen replacement therapy. *Obstet Gynecol Surv.* 1984; 39: 230-245.
8. Beral V, Banks E, Reeves G, Appleby P. Use of HRT and the subsequent risk of cancer. *J Epidemiol Biostat.* 1999; 4: 191-215.
9. Tavani A, LaVecchia C. The adverse effects of hormone replacement therapy. *Drugs Aging.* 1999; 14: 347-357.
10. Perry LM. *Medicinal Plants of East and South East Asia Attributed Properties and Uses*. Cambridge: MIT Press; 1980.
11. Heyne K. *The Useful Indonesian Plants*. Research and Development Agency, The Ministry of Forestry, Jakarta, 1987, 659-703.
12. Goncalves JLS, Lopes RC, Oliveira DB, Costa SS, Miranda MMFS, Romanos MTV et al. *In vitro* antirotavirus activity of some medicinal plants used in Brazil against

- diarrhoea. *J Ethnopharmacol.* 2005; 99: 403-407.
13. Charoenlarp P, Radomyos P, Harinasuta T. Treatment of taeniasis with Puag-Haad: a crude extract of *Artocarpus lakoocha* wood. *The Southeast Asian J Tropical Med Public Health.* 1981; 12: 568-570.
 14. Salguero CP. *A Thai Herbal Traditional Recipes for Health and Harmony*, Scotland, Findhorn Press, 2003, 119.
 15. Joshee N, Bastola DR, Agrawal VP, Yadav AK. Lakoocha: a multipurpose tree of warm climates. In: Janick, J., Whipkey, A. (Eds.), *Trends in New Crops and New Uses*. ASHS Press, Alexandria, VA, 2002.
 16. Likhitwitayawuid K, Sritularak B, Benchanak K, Lipipun V, Mathew J, Schinazi RF. Phenolic with antiviral activity from *Millettia erythrocalyx* and *Artocarpus lakoocha*. *Nat Prod Res.* 2005; 19(2): 177-182.
 17. Saowakon N, Tansatit T, Wanichanon C, Chaakul W, Reutrakul V, Sobhan P. *Fasciola gigantica*: anthelmintic effect of the aqueous extract of *Artocarpus lakoocha*. *Exp Parasitol.* 2009; 122: 289-298.
 18. Tengamnuay P, Pengrungruangwong K, Pheansri I, Likhitwitayawuid K. *Artocarpus lakoocha* heartwood extract as a novel cosmetic ingredient: evaluation of the *in vitro* anti-tyrosinase and *in vivo* skin whitening activities. *Int J Cosmetic Sci.* 2006; 28; 269-276.
 19. The wealth of India "A Dictionary of Indian raw materials and industrial products". 2007; 1: 453-4.
 20. Rahman MA, Hussain Arshad. Anticancer activity and apoptosis inducing effect of methanolic extract of *Cordia dichotoma* against human cancer cell line. *Bangladesh J Pharmacol.* 2015; 10: 27-34.
 21. Trease GE, Evans WC. *A text book of pharmacognosy*. Oxford: ELSB Baillere Tindal; 1987. p. 1055.
 22. Vijayanarayana K, Rodrigues RS, Chandrashekhar KS, Subrahmanyam EV. Evaluation of estrogenic activity of alcoholic extract of rhizome of *Curculigo orchoides*. *J Ethnopharmacol.* 2007; 114: 241-245.
 23. Parhizkar S, Latiff LA, Rahman SA, Dollah MA, Parichehr H. Assessing estrogenic activity of *Nigella sativa* in ovariectomized rats using vaginal cornification assay. *African J Pharm Pharmacol.* 2011; 5: 137-142.
 24. Alam Q, Vijayanarayana K, Satyanarayana D. Evaluation of Estrogenic Activity of Alcoholic Extracts of Fruits of *Solanum xanthocarpum* Using Percentage Vaginal Cornification and Vaginal Opening as Parameters of Assessment. *Pharmacol online.* 2010; 3: 495-502.
 25. Anklesaria BS. The promise of phytoestrogens. *Indian J Pharmacol.* 2011; 43: 369-370.
 26. Hays J, Ockene JK, Brunner RL, Kotchen JM, Manson JE, Patterson RE, Aragaki AK, Shumaker SA, Brzyski RG, LaCroix AZ, Granek IA, Valanis BG. Effects of estrogen plus progestin on health related quality of life. *N Engl J Med.* 2003; 348: 1839-1854.
 27. De Naeyer A, Berghe WV, Pocock V, Milligan S, Haegeman G, De Keukeleire D. Estrogenic and anticarcinogenic properties of kurarinone, a lavandulyl flavanone from the roots of *Sophora flavescens*. *J Nat Prod* 2004; 67: 1829-1832.